LLOYD D LANDSMAN MD

plastic & cosmetic surgery 994 W Jericho Tpke Suite 102 Smithtown NY 11787 631-864-4111 drlandsman.com

Post Operative Instructions

Procedure: Mastopexy/Breast Reduction

- 1. You must spend the first evening following your surgery with an adult.
- 2. Postoperative pain should be moderate. Take the pain medication prescribed as directed. Be sure to take the phenergan with the pain medication in order to minimize postoperative nausea.
- 3. If you should notice any significant swelling, particularly on one side, or if there is persistent oozing of blood or of fluid of any type, please notify the doctor immediately.
- 4. If you should have a fever of greater than 101.6 F, please notify the doctor immediately.
- 5. Drains may be placed at the time of your surgery. Please empty them twice a day. They will be removed at your first postoperative visit.
- 6. Leave your present dressing in place. It will be changed at your first postoperative visit.
- 7. Do not apply any ice or heat to the operative site during the postoperative period.
- 8. No heavy lifting, strenuous exercise or activity is permitted for the first 2 weeks following surgery. After 2 weeks, you may gradually resume your usual level of activity. Any exercise or weight lifting, which relies on the muscles of the chest may be resumed 4 weeks after surgery.
- 9. You may bathe, but try to keep the dressing on the breast area dry at all times.
- 10. Please do not wear any bra except those approved by the doctor. No underwire bras are permitted until 4 weeks following surgery. AT your first post operative visit a post operative bra will be placed following removal of your dressings
- 11. Your first post operative visit will be on _
- 12. If you should have any questions or have any problems please call the office immediately at 631-864-4111, and the doctor will return your call as soon as possible. DO NOT USE EMAIL FOR EMERGENCIES.