

Ali Totonchi, M.D.
(216) 778 - 2262



Care after Abdominoplasty (Tummy Tuck)

This page contains answers to frequently asked questions regarding your care after abdominoplasty. This information will help you to remember the instructions given to you after surgery.

What will I see when I look into a mirror?

It will be difficult to stand up straight for several days. Your abdomen will feel tight and may appear mildly bruised and slightly swollen. The contour of your abdomen will not take its final shape for several weeks after the surgery.

You may have dark discoloration from dry blood at your incision and this is normal. You may notice a “ruffled” appearance to the closure. This is normal and will flatten with time. The belly button incision may develop a yellow crust that is dried fluid. This is normal and will be cleaned at your first post-op visit.

When can I shower or take a bath?

You may not shower until after your first post op visit. Your original dressings and binder from surgery should remain on until your scheduled post op follow up appointment. You may sponge bath but no showers until your drains are removed.

How do I care for my drains?

Your drains are held in place with stitches to your skin and should always be supported to avoid pulling on the drain. The easiest method is to safety pin the drain bulb to your abdominal support or clothing.

The “How to care for your drain” pamphlet has more detailed information. Drains are generally removed within 7 to 10 days after your surgery.

Should I wear a tummy support?

For your comfort, we recommend that you wear an abdominal support garment for 12 weeks following your surgery, removing the garment only for bathing. You can wear the surgical support provided to you for several days-usually until the drains are removed. Most patients find that a spandex or Lycra waist and tummy shaping garment are most comfortable to wear. We will advise you of your garment options before surgery and again at your post-operative

visits. Whatever you choose, it should be easy and comfortable for you to put on and remove. After 6 weeks, you may choose to not wear a support garment for brief periods of time. After 12 weeks, you may discontinue all support garments.

What can I do after surgery?

You will gradually be able to stand upright over the first week. Many patients find it easier to sleep in a recliner for a few days after the surgery. You should walk every day but have someone assist you for the first few days. You should not lift anything heavier than five pounds or participate in heavy exercise for six weeks after the surgery. After six weeks, gradually return to your normal activities and exercise routines. Depending on the type of work you do, you should be able to return to your job as soon as you are comfortable. If your job does not require heavy activity, you should be able to go back in two to three weeks.

When can I drive?

You generally should be able to drive about five to seven days after surgery, if you feel up to it. You should never drive if you are still taking any pain medication other than Tylenol.

How much pain will I have?

The most discomfort that you will have lasts three to five days, sometimes a little longer. As you feel more comfortable, your need for medication will lessen.

What medications should I use?

Your surgeon will prescribe a pain medication for you and may also have you take an antibiotic. Follow the directions on the bottle for their use. The pain medication will make you feel drowsy. Have someone assist you in your home and do not attempt to drive while you are taking the pain medication. A few days after surgery most patients will find that Tylenol will take care of the discomfort. Do not use aspirin or ibuprofen (Motrin) for five days after the surgery.

Can I drink alcohol?

Do not drink alcohol for five days after your surgery or while taking narcotic pain medicines.

When will I be seen in the office after surgery?

You will normally be seen in the office within one week of your surgery and then at about two weeks, six weeks, and six months after surgery to monitor healing. Additional appointments may be necessary.

How will I know if I am having a problem?

Urgent problems after abdominoplasty do not occur frequently. These are the complications that will require a change in your post-operative care:

Hematoma (blood collecting under your incisions) can occur within a few days of your surgery. The warning signs are:

- Severe pain that does not respond to medication
- Significant swelling in your abdomen
- Excessive or growing bruising

Infection is rare following abdominoplasty surgery. It is normal to have a small amount of drainage from the incisions and around your belly button for a few days. Signs of infection are:

- Increased temperature
- Increasing drainage from the incisions
- Increasing redness around the incisions

Leg swelling with or without associated pain may indicate a problem with the vein circulation in your leg(s). Although some swelling is to be expected due to your surgery, intravenous fluids given to you during the surgery, and your decreased level of activity, this swelling is normally mild, painless and affects both legs evenly. If you have a large amount of leg swelling (either one or both legs) or if you experience pain in your legs, contact the office immediately.

Breathing problems after abdominoplasty are rare but can be a serious complication. If you develop any chest and/or back pain or the feeling of being short of breath you must contact my office or be seen in the nearest emergency medical facility without delay.

Medication reactions may occur with the drugs prescribed for you. If you develop a skin rash, itching, vomiting, or diarrhea, stop taking your medication and contact my office.

If you have any of the above problems or other questions, contact my office, 216-778-4450.