

Ali Totonchi, M. D.

Care after Frontal Bossing

This page contains answers to frequently asked questions regarding your care after Frontal Bossing surgery. This information will assist you to remember the instructions given to you after surgery.

What will I see when I look in the mirror?

Swelling and bruising are to be expected. You may have bruising and swelling around the eyes.

Keep all dressings that have been placed on you at the time of surgery intact until you are seen at your first post-operative appointment. The dressings stay in place for 3-5 days. Please keep the dressings clean and dry.

The incision will be repaired with very fine sutures in the hairline. These sutures will dissolve on their own. Once the dressings are removed in the office, you will begin applying bacitracin ointment to the incision twice daily for 1 week. Remove old bacitracin before applying new bacitracin.

When can I shower or take a bath?

You can bathe normally, be sure to keep any dressings dry.

Do not shower until dressings have been removed at your post op visit.

What can I do after surgery?

You should take it easy for several days after surgery and gradually increase your activity as you feel more comfortable.

You should sleep in a semi-reclined position for 1 week after surgery to minimize swelling.

You should NOT do any heavy lifting, greater than 5-10 lb, or strenuous exercise for 3 weeks. You can restart your exercise regime gradually, as tolerated, 3 weeks after your surgery.

When can I drive?

You will not be able to drive yourself if you are taking narcotic pain medication.

How much pain will I have after surgery?

The most discomfort you will have lasts the first 3-5 days after surgery.

You will be given a prescription for pain medication, take it as directed, if needed for pain. The medication may make you feel drowsy so please have someone assist you at home for the first few days after surgery.

Do not take any Aspirin, Advil, Motrin, or Aleve for 3 weeks before surgery and avoid for 1 week after surgery.

When can I drink alcohol?

You can drink alcohol 7 days after your procedure if you are no longer taking pain medication.

When will I be seen in the office after surgery?

You will normally be seen in the office 3-5 days after surgery to monitor and remove the dressings and then at 1 month and 6 months to monitor healing.

How will I know if I am having a problem?

Hematoma (collection of blood under the skin) can occur within a few hours to a few days of your surgery.

The warning signs are:

- Severe pain that does not respond to medication
- Significant swelling in an isolated area
- Excessive bruising in one area compared to another

Infection Signs of infection are:

- Increased redness around the incisions
- Increased drainage from the incisions
- Increased temperature/fever

Leg swelling (with or without associated pain) may indicate a problem with the vein circulation in your leg(s). If you have a large amount of leg swelling (either in one or both legs), or if you experience pain in your legs, contact the office immediately.

Breathing problems with or without chest pain, seek medical attention immediately.

Medication reactions may occur with the drugs prescribed for you. If you develop a skin rash, itching, vomiting, or diarrhea, or other concerns, stop taking your medication and call the office.

If you have any of the above problems or other questions, contact my office at 440-461-7999.